

An abstract painting with a complex, layered composition. The background is a mix of light and dark tones, with prominent splatters and brushstrokes in vibrant colors: bright blue, deep red, yellow, and green. There are also some darker, more muted tones like purple and black. The overall effect is one of intense energy and emotional turbulence, with various shapes and textures that suggest movement and depth.

When Worry Gets Out of Control:  
**Generalized Anxiety  
Disorder**

**Easy To Read**

**National Institute of Mental Health**

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES • National Institutes of Health



# Generalized Anxiety Disorder

## Anxiety Disorders

People with anxiety disorders feel extremely fearful and unsure. Most people feel anxious about something for a short time now and again, but people with anxiety disorders feel this way most of the time. Their fears and worries make it hard for them to do everyday tasks. About 18% of American adults have anxiety disorders. Children also may have them.

Treatment is available for people with anxiety disorders. Researchers are also looking for new treatments that will help relieve symptoms.

This booklet is about one kind of anxiety disorder called generalized anxiety disorder, or GAD. For information about other kinds of anxiety disorders, please see the end of this booklet.



## **Generalized Anxiety Disorder**

All of us worry about things like health, money, or family problems at one time or another. But people with GAD are extremely worried about these and many other things, even when there is little or no reason to worry about them. They may be very anxious about just getting through the day. They think things will always go badly. At times, worrying keeps people with GAD from doing everyday tasks.

People with GAD may be very  
anxious about just getting  
through the day. They think things  
will always go badly.

## This is a list of common symptoms.

### People with GAD:

- **worry very much about everyday things** for at least six months, even if there is little or no reason to worry about them;
- **can't control** their constant worries;
- **know that they worry** much more than they should;
- **can't relax**;
- **have a hard time** concentrating;
- **are easily startled**; and
- **have trouble** falling asleep or staying asleep.
- **common body symptoms are:**
  - feeling tired for no reason;
  - headaches;
  - muscle tension and aches;
  - having a hard time swallowing;
  - trembling or twitching;
  - being irritable;
  - sweating;
  - nausea;
  - feeling lightheaded;
  - feeling out of breath;
  - having to go to the bathroom a lot; and
  - hot flashes.

## **When does GAD start?**

GAD develops slowly. It often starts during the time between childhood and middle age. Symptoms may get better or worse at different times, and often are worse during times of stress.

People with GAD may visit a doctor many times before they find out they have this disorder. They ask their doctors to help them with the signs of GAD, such as headaches or trouble falling asleep, but don't always get the help they need right away. It may take doctors some time to be sure that a person has GAD instead of something else.

**There is help for people with GAD.**

**The first step is to go to a doctor to talk about symptoms.**

## Is there help?

There is help for people with GAD. The first step is to go to a doctor or health clinic to talk about symptoms. People who think they have GAD may want to bring this booklet to the doctor, to help them talk about the symptoms in it. The doctor will do an exam to make sure that another physical problem isn't causing the symptoms. The doctor may make a referral to a mental health specialist.

Doctors may prescribe medication to help relieve GAD. It's important to know that some of these medicines may take a few weeks to start working. In most states only a medical doctor (a family doctor or psychiatrist) can prescribe medications.

The kinds of medicines used to treat GAD are listed below. Some are used to treat other problems, such as depression, but also are helpful for GAD:

- antidepressants,
- anti-anxiety medicines, and
- beta blockers.

Doctors also may ask people with GAD to go to therapy with a licensed social worker, psychologist, or psychiatrist. This treatment can help people with GAD feel less anxious and fearful.

There is no cure for GAD yet, but treatments can give relief to people who have it and help them live a more normal life. If you know someone with signs of GAD, talk to him or her about seeing a doctor. Offer to go along for support.

**To find out more about GAD, call 1-866-615-NIMH (1-866-615-6464) to have free information mailed to you.**

## **Who pays for treatment?**

Most insurance plans cover treatment for anxiety disorders. People who are going to have treatment should check with their own insurance companies to find out about coverage. For people who don't have insurance, local city or county governments may offer treatment at a clinic or health center, where the cost is based on income. Medicaid plans also may pay for GAD treatment.

## **Why do people get GAD?**

GAD sometimes runs in families, but no one knows for sure why some people have it, while others don't. When chemicals in the brain are not at a certain level it can cause a person to have GAD. That is why medications often help with the symptoms because they help the brain chemicals stay at the correct levels.

To improve treatment, scientists are studying how well different medicines and therapies work. In one kind of research, people with GAD choose to take part in a clinical trial to help doctors find out what treatments work best for most people, or what works best for different symptoms. Usually, the treatment is free. Scientists are learning more about how the brain works so that they can discover new treatments.





## Personal story

**“I was worried all the time about everything. It didn’t matter that there were no signs of problems, I just got upset. I was having trouble falling asleep at night, and I couldn’t keep my mind focused at work. I felt angry at my family all the time.”**

**“I saw my doctor and explained my constant worries. My doctor sent me to someone who knows about GAD. Now I am taking medicine and working with a counselor to cope better with my worries. I had to work hard, but I feel better. I’m glad I made that first call to my doctor.”**

## **For More Information on GAD and Other Anxiety Disorders**

Visit the National Library of Medicine's  
MedlinePlus [www.nlm.nih.gov/medlineplus](http://www.nlm.nih.gov/medlineplus)  
En Español, <http://medlineplus.gov/spanish>

## **For Information on Clinical Trials for GAD and Other Anxiety Disorders**

NIMH Clinical Trials Web page  
[www.nimh.nih.gov/studies/index.cfm](http://www.nimh.nih.gov/studies/index.cfm)

National Library of Medicine Clinical Trials Database  
[www.clinicaltrials.gov](http://www.clinicaltrials.gov)

Information from NIMH is available in multiple formats.  
You can browse online, download documents in PDF,  
and order paper brochures by mail. You can order NIMH  
publications online at [www.nimh.nih.gov](http://www.nimh.nih.gov). If you do not  
have Internet access, please contact the NIMH Information  
Center at the numbers listed below.

## **For Further Information**

### **National Institute of Mental Health**

Science Writing, Press, & Dissemination Branch  
6001 Executive Boulevard  
Room 8184, MSC 9663  
Bethesda, MD 20892-9663  
Phone: 301-443-4513 or  
1-866-615-NIMH (6464) toll-free  
TTY: 301-443-8431  
TTY: 866-415-8051 toll-free  
FAX: 301-443-4279  
E-mail: [nimhinfo@nih.gov](mailto:nimhinfo@nih.gov)  
Web site: [www.nimh.nih.gov](http://www.nimh.nih.gov)

This publication is in the public domain and may be reproduced or copied without permission from the National Institute of Mental Health (NIMH). NIMH encourages you to reproduce this publication and use it in your efforts to improve public health. Citation of the NIMH as a source is appreciated. However, using government materials inappropriately can raise legal or ethical concerns, so we ask you to use these guidelines:

- NIMH does not endorse or recommend any commercial products, processes, or services, and this publication may not be used for advertising or endorsement purposes.
- NIMH does not provide specific medical advice or treatment recommendations or referrals; these materials may not be used in a manner that has the appearance of such information.
- NIMH requests that Non-Federal organizations not alter this publication in a way that will jeopardize the integrity and “brand” when using the publication.
- Addition of Non-Federal Government logos and Web site links may not have the appearance of NIMH endorsement of any specific commercial products or services or medical treatments or services.

If you have questions regarding these guidelines and use of NIMH publications, please contact the NIMH Information Center at 1-866-615-6464 or e-mail at [nimhinfo@nih.gov](mailto:nimhinfo@nih.gov).



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES  
National Institutes of Health  
NIH Publication No. 07-4677  
Revised 2007